


## AMENDMENTS TO THE CLAIMS

### IN THE CLAIMS:

This listing of claims will replace all prior versions and listings of claims in the application. Please amend the claims as follows.

1. (Currently amended) A compound of Formula (I):



wherein

m is an integer of from 0 to 3;

n is an integer of from 0 to 3;


R<sub>1</sub> is aryl;

R<sub>2</sub> is

- a) a group of the formula -N(R<sub>9</sub>R<sub>10</sub>), -NHC(O)R<sub>9</sub>, or -NHC(O)OR<sub>9</sub>;
- ~~b) a group of the formula -OR<sub>9</sub>;~~
- ~~c) a group of the formula -SR<sub>9</sub>, -SOR<sub>9</sub>, -SO<sub>2</sub>R<sub>9</sub>, -SO<sub>2</sub>NHR<sub>9</sub>, or -SO<sub>2</sub>N(R<sub>9</sub>R<sub>10</sub>);~~

wherein R<sub>9</sub> and R<sub>10</sub> are independently selected from the group consisting of

- 1) -H;
- 2) -Aryl;
- 3) -C<sub>1-6</sub> alkyl;
- 4) -C<sub>1-6</sub> alkylaryl;



6) -aryl; and

7) -C<sub>1-6</sub> alkyl;

R<sub>3</sub> and R<sub>4</sub> are independently selected from the group consisting of


- a) H;
- b) -aryl;
- c) -C<sub>1-6</sub> alkyl;
- d) -C<sub>1-6</sub> alkylaryl; and
- e) -C<sub>1-6</sub> alkoxyaryl;

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> are independently selected from the group consisting of

- a) -H;
- b) -C<sub>1-6</sub> alkyl;
- c) -aryl;
- d) -C<sub>1-6</sub> alkylaryl;
- e) -C(O)-O-C<sub>1-6</sub> alkyl;
- f) -C(O)-O-C<sub>1-6</sub> alkylaryl;
- g) -C(O)-NH-C<sub>1-6</sub> alkyl;
- h) -C(O)-NH-C<sub>1-6</sub> alkylaryl;
- i) -SO<sub>2</sub>-C<sub>1-6</sub> alkyl;
- j) -SO<sub>2</sub>-C<sub>1-6</sub> alkylaryl;
- k) -SO<sub>2</sub>-aryl;
- l) -SO<sub>2</sub>-NH-C<sub>1-6</sub> alkyl;
- m) -SO<sub>2</sub>-NH-C<sub>1-6</sub> alkylaryl;
- n) -C(O)-C<sub>1-6</sub> alkyl;
- o) -C(O)-C<sub>1-6</sub> alkylaryl;

- p) -Y-C<sub>1-6</sub> alkyl;
- q) -Y-aryl;
- r) -Y-C<sub>1-6</sub> alkylaryl;
- s) -Y-C<sub>1-6</sub> alkylene-NR<sub>13</sub>R<sub>14</sub>;
- t) -Y-C<sub>1-6</sub> alkylene-W-R<sub>15</sub>;

wherein Y and W are independently selected from the group consisting of -CH<sub>2</sub>-, -O-, -N(H)-, -S-, SO<sub>2</sub>-, -CON(H)-, -NHC(O)-, -NHCON(H)-, -NHSO<sub>2</sub>-, -SO<sub>2</sub>N(H)-, -C(O)-O-, -NHSO<sub>2</sub>NH-, -O-CO-,



wherein R<sub>16</sub> and R<sub>17</sub> are independently selected from the group consisting of aryl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkylaryl, C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>1</sub>-C<sub>6</sub> alkoxyaryl;

R<sub>15</sub> is aryl, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>6</sub> alkylaryl; and


- u) halogen, hydroxyl, cyano, carbamoyl, and carboxyl;

wherein at least one of R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> is -Y-C<sub>1-6</sub> alkylene-NR<sub>13</sub>R<sub>14</sub>, and

R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, and R<sub>14</sub> are independently selected from the group consisting of hydrogen, aryl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkylaryl, C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>1</sub>-C<sub>6</sub> alkoxyaryl; or

R<sub>13</sub> and R<sub>14</sub> are taken together to form a ring having the formula -(CH<sub>2</sub>)<sub>o</sub>-X-(CH<sub>2</sub>)<sub>p</sub>- bonded to the nitrogen atom to which R<sub>13</sub> and R<sub>14</sub> are attached, and/or R<sub>11</sub> and R<sub>12</sub> are taken together to form a ring having the formula -(CH<sub>2</sub>)<sub>o</sub>-X-(CH<sub>2</sub>)<sub>p</sub>- bonded to the atoms to which R<sub>11</sub> and R<sub>12</sub> are connected, wherein o and p are, independently, 1, 2, 3, or 4; X


is a direct bond,  $-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{S}(\text{O}_2)-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{CON}(\text{H})-$ ,  $-\text{NHC}(\text{O})-$ ,  $-\text{NHCON}(\text{H})-$ ,  $-\text{NHSO}_2-$ ,  $-\text{SO}_2\text{N}(\text{H})-$ ,  $-\text{C}(\text{O})-\text{O}-$ ,  $-\text{O}-\text{C}(\text{O})-$ ,  $-\text{NHSO}_2\text{NH}-$ ,



wherein the aryl and/or alkyl group(s) in  $\text{R}_1$ ,  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_5$ ,  $\text{R}_6$ ,  $\text{R}_7$ ,  $\text{R}_8$ ,  $\text{R}_9$ ,  $\text{R}_{10}$ ,  $\text{R}_{11}$ ,  $\text{R}_{12}$ ,  $\text{R}_{13}$ ,  $\text{R}_{14}$ ,  $\text{R}_{15}$ ,  $\text{R}_{16}$ ,  $\text{R}_{17}$ ,  $\text{R}_{18}$ , and  $\text{R}_{19}$  may be optionally substituted 1-4 times with a substituent group, wherein said substituent group(s) or the term substituted refers to groups selected from the group consisting of:

- a)  $-\text{H}$ ;
- b)  $-\text{Z}-\text{C}_{1-6}$  alkyl;
- $-\text{Z}-\text{aryl}$ ;
- $-\text{Z}-\text{C}_{1-6}$  alkylaryl;
- $-\text{Z}-\text{C}_{1-6}-\text{alkyl}-\text{NR}_{20}\text{R}_{21}$ ;
- $-\text{Z}-\text{C}_{1-6}-\text{alkyl}-\text{W}-\text{R}_{22}$ ;


wherein Z and W are independently selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{N}(\text{H})-$ ,  $-\text{S}-$ ,  $\text{SO}_2-$ ,  $-\text{CON}(\text{H})-$ ,  $-\text{NHC}(\text{O})-$ ,  $-\text{NHCON}(\text{H})-$ ,  $-\text{NHSO}_2-$ ,  $-\text{SO}_2\text{N}(\text{H})-$ ,  $-\text{C}(\text{O})-\text{O}-$ ,  $-\text{NHSO}_2\text{NH}-$ ,  $-\text{O}-\text{CO}-$ ,



$R_{22}$ ,  $R_{23}$ , and  $R_{24}$  are independently selected from the group consisting of aryl,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkylaryl,  $C_1$ - $C_6$  alkoxy, and  $C_1$ - $C_6$  alkoxyaryl;


c) halogen, hydroxyl, cyano, and carbamoyl; and

wherein  $R_{20}$  and  $R_{21}$  are independently selected from the group consisting of hydrogen, aryl,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkylaryl,  $C_1$ - $C_6$  alkoxy, and  $C_1$ - $C_6$  alkoxyaryl; or  $R_{20}$  and  $R_{21}$  are taken together to form a ring having the formula  $-(CH_2)_q-X-(CH_2)_r-$  bonded to the nitrogen atom to which  $R_{20}$  and  $R_{21}$  are attached wherein  $q$  and  $r$  are, independently, 1, 2, 3, or 4;  $X$  is a direct bond,  $-CH_2-$ ,  $-O-$ ,  $-S-$ ,  $-S(O_2)-$ ,  $-C(O)-$ ,  $-CON(H)-$ ,  $-NHC(O)-$ ,  $-NHCON(H)-$ ,  $-NHSO_2-$ ,  $-SO_2N(H)-$ ,  $-C(O)-O-$ ,  $-O-C(O)-$ ,  $-NHSO_2NH-$ ,



$R_{25}$ ,  $R_{26}$ , and  $R_{27}$  are independently selected from the group consisting of hydrogen, aryl,  $C_1$ - $C_6$  alkyl, and  $C_1$ - $C_6$  alkylaryl; or a pharmaceutically acceptable salt, solvate or prodrug thereof.

2. (Previously presented) The compound of claim 1, wherein  $m$  is an integer of from 0 to 3;  
 $n$  is 0;  $R_3$  is hydrogen as represented by the formula (II)



and wherein

$R_1$  is an aryl group;

$R_2$  is a group of the formula  $-N(R_9R_{10})$ ,  $-NHC(O)R_9$ , or  $-NHC(O)OR_9$ ;

wherein  $R_9$  and  $R_{10}$  are independently selected from the group consisting of

- 1)  $-H$ ;
- 2)  $-Aryl$ ;
- 3)  $-C_{1-6}$  alkyl; and
- 4)  $-C_{1-6}$  alkylaryl;

$R_4$  is


- a)  $H$ ;
- b)  $-aryl$ ;
- c)  $-C_{1-6}$  alkyl;
- d)  $-C_{1-6}$  alkylaryl; or
- e)  $-C_{1-6}$  alkoxyaryl;

$R_5$ ,  $R_6$ ,  $R_7$ , and  $R_8$  are independently selected from the group consisting of

- a)  $-H$ ;
- b)  $-C_{1-6}$  alkyl;
- c)  $-aryl$ ;
- d)  $-C_{1-6}$  alkylaryl;
- e)  $-C(O)-O-C_{1-6}$  alkyl;

- f) -C(O)-O-C<sub>1-6</sub> alkylaryl;
- g) -C(O)-NH-C<sub>1-6</sub> alkyl;
- h) -C(O)-NH-C<sub>1-6</sub> alkylaryl;
- i) -SO<sub>2</sub>-C<sub>1-6</sub> alkyl;
- j) -SO<sub>2</sub>-C<sub>1-6</sub> alkylaryl;
- k) -SO<sub>2</sub>-aryl;
- l) -SO<sub>2</sub>-NH-C<sub>1-6</sub> alkyl;
- m) -SO<sub>2</sub>-NH-C<sub>1-6</sub> alkylaryl
- n) -C(O)-C<sub>1-6</sub> alkyl;
- o) -C(O)-C<sub>1-6</sub> alkylaryl;
- p) -Y-C<sub>1-6</sub> alkyl;
- q) -Y-aryl;
- r) -Y-C<sub>1-6</sub> alkylaryl;
- s) -Y-C<sub>1-6</sub> alkylene-NR<sub>13</sub>R<sub>14</sub>;
- t) -Y-C<sub>1-6</sub> alkylene-W-R<sub>15</sub>;

wherein Y and W are independently selected from the group consisting of -CH<sub>2</sub>-, -O-, -N(H)-, -S-, SO<sub>2</sub>-, -CON(H)-, -NHC(O)-, -NHCON(H)-, -NHSO<sub>2</sub>-, -SO<sub>2</sub>N(H)-, -C(O)-O-, -NHSO<sub>2</sub>NH-, -O-CO-,



wherein R<sub>16</sub> and R<sub>17</sub> are independently selected from the group consisting of aryl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkylaryl, C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>1</sub>-C<sub>6</sub> alkoxyaryl;


R<sub>15</sub> is aryl, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>6</sub> alkylaryl, and

u) halogen, hydroxyl, cyano, carbamoyl, and carboxyl;

wherein at least one of R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> is -Y-C<sub>1-6</sub> alkylene-N-R<sub>13</sub>R<sub>14</sub>,

R<sub>13</sub>, and R<sub>14</sub> are independently selected from the group consisting of hydrogen, aryl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkylaryl, C<sub>1</sub>-C<sub>6</sub> alkoxy, and C<sub>1</sub>-C<sub>6</sub> alkoxyaryl; or

R<sub>13</sub> and R<sub>14</sub> are together to form a ring having the formula -(CH<sub>2</sub>)<sub>o</sub>-X-(CH<sub>2</sub>)<sub>p</sub>- bonded to the nitrogen atom to which R<sub>13</sub> and R<sub>14</sub> are attached, wherein o and p are, independently, 1, 2, 3, or 4; X is a direct bond, -CH<sub>2</sub>-, -O-, -S-, -S(O<sub>2</sub>)-, -C(O)-, -CON(H)-, -NHC(O)-, -NHCON(H)-, -NHSO<sub>2</sub>-, -SO<sub>2</sub>N(H)-, -C(O)-O-, -O-C(O)-, -NHSO<sub>2</sub>NH-,




and wherein the aryl and/or alkyl group(s) in R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub>, R<sub>18</sub>, and R<sub>19</sub> may be optionally substituted 1-4 times with a substituent group, wherein said substituent group(s) or the term substituted refers to groups selected from the group consisting of:

- a) -H;
- b) -Z-C<sub>1-6</sub> alkyl;  
 -Z-aryl;  
 -Z-C<sub>1-6</sub> alkylaryl;  
 -Z-C<sub>1-6</sub>-alkyl-NR<sub>20</sub>R<sub>21</sub>;  
 -Z-C<sub>1-6</sub>-alkyl-W-R<sub>22</sub>;



wherein Z and W are independently selected from the group consisting of  $-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{N}(\text{H})-$ ,  $-\text{S}-$ ,  $-\text{SO}_2-$ ,  $-\text{CON}(\text{H})-$ ,  $-\text{NHC}(\text{O})-$ ,  $-\text{NHCON}(\text{H})-$ ,  $-\text{NHSO}_2-$ ,  $-\text{SO}_2\text{N}(\text{H})-$ ,  $-\text{C}(\text{O})-\text{O}-$ ,  $-\text{NHSO}_2\text{NH}-$ ,  $-\text{O}-\text{CO}-$ ,




wherein;

$\text{R}_{22}$ ,  $\text{R}_{23}$ , and  $\text{R}_{24}$  are independently selected from the group consisting of aryl,  $\text{C}_1$ - $\text{C}_6$  alkyl,  $\text{C}_1$ - $\text{C}_6$  alkylaryl,  $\text{C}_1$ - $\text{C}_6$  alkoxy, and  $\text{C}_1$ - $\text{C}_6$  alkoxyaryl;

c) halogen, hydroxyl, cyano, and carbamoyl; and

wherein  $\text{R}_{20}$  and  $\text{R}_{21}$  are independently selected from the group consisting of hydrogen, aryl,  $\text{C}_1$ - $\text{C}_6$  alkyl,  $\text{C}_1$ - $\text{C}_6$  alkylaryl,  $\text{C}_1$ - $\text{C}_6$  alkoxy, and  $\text{C}_1$ - $\text{C}_6$  alkoxyaryl; or

$\text{R}_{20}$  and  $\text{R}_{21}$  are taken together to form a ring having the formula  $-(\text{CH}_2)_q-\text{X}-(\text{CH}_2)_r-$  bonded to the nitrogen atom to which  $\text{R}_{20}$  and  $\text{R}_{21}$  are attached wherein q and r are, independently, 1, 2, 3, or 4; X is a direct bond,  $-\text{CH}_2-$ ,  $-\text{O}-$ ,  $-\text{S}-$ ,  $-\text{S}(\text{O}_2)-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{CON}(\text{H})-$ ,  $-\text{NHC}(\text{O})-$ ,  $-\text{NHCON}(\text{H})-$ ,  $-\text{NHSO}_2-$ ,  $-\text{SO}_2\text{N}(\text{H})-$ ,  $-\text{C}(\text{O})-\text{O}-$ ,  $-\text{O}-\text{C}(\text{O})-$ ,  $-\text{NHSO}_2\text{NH}-$ ,



R<sub>25</sub> and R<sub>26</sub> are independently selected from the group consisting of hydrogen, aryl, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>1</sub>-C<sub>6</sub> alkylaryl; or a pharmaceutically acceptable salt, solvate or prodrug thereof.

3. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-tert-butoxycarbonylamino-1-ethyl]-3-butyl-5-(3-diethylamino-1-propoxy)benzimidazole.

4. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-amino-1-ethyl]-3-butyl-5-(3-diethylamino-1-propoxy)benzimidazole Trihydrochloride.

5. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-tert-butoxycarbonylamino-1-ethyl]-3-butyl-6-(3-diethylamino-1-propoxy)benzimidazole.

6. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-amino-1-ethyl]-3-butyl-6-(3-diethylamino-1-propoxy)benzimidazole.

7. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-tert-butoxycarbonylamino-1-ethyl]-6-(3-diethylamino-1-propoxy)benzimidazole.

8. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-amino-1-ethyl]-6-(3-diethylamino-1-propoxy)benzimidazole.

9. (Previously presented) The compound of claim 1, wherein the compound is 2-[2-(3-Benzyloxyphenyl)-1-(tert-butoxycarbonylamino)-1-ethyl]-3-butyl-5-(3-diethylamino-1-propoxy)benzimidazole.

10. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Ethoxyphenyl)-1-(tert-butoxycarbonylamino)-1-ethyl]-3-butyl-5-(3-diethylamino-1-propoxy)benzimidazole.

11. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-(4-Chloro)phenethoxy)phenyl)-1-(tert-butoxycarbonylamino)-1-ethyl]-3-butyl-5-(3-diethylamino-1-propoxy)benzimidazole.

12. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-(tert-butoxycarbonylamino)-1-ethyl]-3-(3-diethylamino-1-propyl)-5-(3-diethylamino-1-propoxy)benzimidazole.

13. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-(tert-butoxycarbonylamino)-1-ethyl]-3-ethyl-5-(3-diethylamino-1-propoxy)benzimidazole.

14. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-amino-1-ethyl]-3-(3-diethylamino-1-propyl)-5-(3-diethylamino-1-propoxy)benzimidazole.

15. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-(tert-butoxycarbonylamino)-1-ethyl]-3-benzyl-5-(3-diethylamino-1-propoxy)benzimidazole.

16. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-amino-1-ethyl]-3-benzyl-5-(3-diethylamino-1-propoxy)benzimidazole.

17. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-(tert-butoxycarbonylamino)-1-ethyl]-3-propyl-5-(3-diethylamino-1-propoxy)benzimidazole

18. (Previously presented) The compound of claim 1, wherein the compound is 2-[(1R)-2-(4-Benzyloxyphenyl)-1-amino-1-ethyl]-3-propyl-5-(3-diethylamino-1-propoxy)benzimidazole.

19. (Original) A pharmaceutical composition comprising the compound of Formula (I) as claimed in claim 1, and one or more pharmaceutically acceptable carriers, excipients, or diluents.

20. (Currently amended) The pharmaceutical composition of claim 19, in the form of an oral dosage or parenteral dosage unit.

21. (Original) The pharmaceutical composition of claim 19, wherein said compound is administered as a dose in a range from about 0.01 to 500 mg/kg of body weight per day.

22. (Original) The pharmaceutical composition of claim 19, wherein said compound is administered as a dose in a range from about 0.1 to 200 mg/kg of body weight per day.

23. (Original) The pharmaceutical composition of claim 19, wherein said compound is administered as a dose in a range from about 0.1 to 100 mg/kg of body weight per day.

24. (Original) The pharmaceutical composition of claim 19, further comprising one or more therapeutic agents selected from the group consisting of alkylating agents, antimetabolites, plant alkaloids, antibiotics, hormones, biologic response modifiers, analgesics, NSAIDs, DMARDs, glucocorticoids, sulfonyleureas, biguanides, insulin, cholinesterase inhibitors, antipsychotics, antidepressants, and anticonvulsants.

25. (Withdrawn) A method for the inhibition of the interaction of RAGE with its physiological ligands, which comprises administering to a subject in need thereof, at least one compound of Formula (I) as claimed in claim 1.

26. (Withdrawn) The method of claim 25, wherein the ligand(s) is(are) selected from advanced glycated end products (AGEs), S100/calgranulin/EN-RAGE,  $\beta$ -amyloid and amphotericin.

27. (Withdrawn) A method for treating a disease state selected from the group consisting of acute and chronic inflammation, symptoms of diabetes, vascular permeability, nephropathy, atherosclerosis, retinopathy, Alzheimer's disease, erectile dysfunction, and tumor invasion and/or metastasis, which comprises administering to a subject in need thereof a therapeutically effective amount of at least one compound of Formula (I) as claimed in claim 1.

28. (Withdrawn –currently amended) A method ~~of prevention and/or treatment of treating~~ RAGE mediated human diseases comprising administration to a human in need thereof a therapeutically effective amount of a compound of Formula (I) as claimed

in claim 1, wherein a therapeutically effective amount comprises sufficient compound to at least partially inhibit the binding of a physiological ligand to the RAGE receptor.

29. (Withdrawn) The method of claim 28, further comprising administering to a subject in need thereof at least one adjuvant and/or additional therapeutic agent(s).

30. (Withdrawn) The method of claim 29, wherein said therapeutic agents are selected from the group consisting of alkylating agents, antimetabolites, plant alkaloids, antibiotics, hormones, biologic response modifiers, analgesics, NSAIDs, DMARDs, glucocorticoids, sulfonylureas, biguanides, insulin, cholinesterase inhibitors, antipsychotics, antidepressants, and anticonvulsants.

31. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises acute and/or chronic inflammation.

32. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises abnormal vascular permeability.

33. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises nephropathy.

34. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises atherosclerosis.

35. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises retinopathy.

36. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises Alzheimer's disease.

37. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises erectile dysfunction.

38. (Withdrawn) The method of claim 28, wherein the RAGE mediated human disease comprises tumor invasion and/or metastasis.

39. (Previously presented) The compound of claim 1, wherein R<sub>4</sub> is

- a) -aryl;
- b) -C<sub>1-6</sub> alkyl;
- c) -C<sub>1-6</sub> alkylaryl; or
- d) -C<sub>1-6</sub> alkoxyaryl.